

Bethe-Salpeter eigenvalues and amplitudes for the half-filled two-dimensional Hubbard model

N. Bulut

Institute for Theoretical Physics, University of California, Santa Barbara, California 93106-4030

D. J. Scalapino

Department of Physics, University of California, Santa Barbara, California 93106-9530

S. R. White

Department of Physics, University of California, Irvine, California 92717

(Received 19 October 1992)

Monte Carlo simulations are used to determine the eigenvalues and eigenfunctions of the particle-hole and particle-particle Bethe-Salpeter equations for 8×8 half-filled Hubbard lattice with $U/t=4$ and $U/t=8$. In the particle-hole channel, the dominant eigenvalue corresponds to the $\mathbf{Q}=(\pi, \pi)$ antiferromagnetic correlations. In the particle-particle channel the amplitude of the leading low-temperature eigenvalue is an even-frequency $d_{x^2-y^2}$ singlet. Odd-frequency p -wave-singlet and s -wave-triplet amplitudes are also found.

The nature and interplay of the antiferromagnetic and pairing correlations in the two-dimensional Hubbard model near half-filling remain open questions.¹ In the Hubbard model, the electrons responsible for the antiferromagnetic fluctuations are also the electrons that pair. Thus one would like to examine the particle-hole spin fluctuation and the particle-particle pairing channels on an equal footing. In order to do this we have carried out Monte Carlo calculations of the two-fermion scattering vertex on an 8×8 half-filled two-dimensional Hubbard lattice with

$$H = -t \sum_{\langle i,j \rangle, \sigma} (c_{i\sigma}^\dagger c_{j\sigma} + c_{j\sigma}^\dagger c_{i\sigma}) + U \sum_i (n_{i\uparrow} - \frac{1}{2})(n_{i\downarrow} - \frac{1}{2}). \quad (1)$$

Here $c_{i\sigma}^\dagger$ creates an electron of spin σ on site i , t is a near-neighbor hopping amplitude, and U is the onsite Coulomb repulsion. Equation (1) is written in a particle-hole symmetric form appropriate for a half-filled band. Viewed in the particle-hole channel with center-of-mass momentum $\mathbf{Q}=(\pi, \pi)$, the two-fermion vertex provides direct information on the antiferromagnetic correlations. Viewed in the $\mathbf{q}=(0,0)$ center-of-mass particle-particle channel, it provides information on the nature of the pairing correlations. Here, combining Monte Carlo results for the two-fermion vertex with Monte Carlo calculations of the single-fermion Green's function, we determine the eigenvalues and amplitudes for the $\mathbf{Q}=(\pi, \pi)$ particle-hole and the $\mathbf{q}=(0,0)$ particle-particle Bethe-

Salpeter equations. As the temperature is lowered, the dominant eigenvalue, which approaches 1 as T goes to zero, occurs in the particle-hole channel. This signals the formation of the zero-temperature antiferromagnetic phase of the half-filled system. In the particle-particle channel, which describes intermediate states doped with two additional fermions,² the dominant eigenvalue at low temperatures is associated with a singlet $d_{x^2-y^2}$ even-frequency amplitude. Additional eigenvalues with odd-frequency pairing amplitudes are also found.

At half-filling, there are no fermion determinantal sign problems,³ and it is straightforward to calculate the one- and two-fermion propagators,

$$G(x_2, x_1) = -\langle T_\tau c_\sigma(x_2) c_\sigma^\dagger(x_1) \rangle \quad (2)$$

and

$$G_2(x_4, x_3, x_2, x_1) = -\langle T_\tau c_{\sigma_4}(x_4) c_{\sigma_3}(x_3) c_{\sigma_2}^\dagger(x_2) c_{\sigma_1}^\dagger(x_1) \rangle. \quad (3)$$

Here $x_l=(l, \tau_l)$ and T_τ is the usual τ -ordering operator. Fourier transforming on both the space- and imaginary-time variables provides information in momentum \mathbf{p} and Matsubara frequency $\omega_n=(2n+1)\pi T$. From the single-particle propagator, Eq. (2), one obtains $G(p)$, and from the two-particle propagator, Eq. (3), $G_2(p_4, p_3, p_2, p_1)$. Here p stands for $(\mathbf{p}, i\omega_n)$ and σ .

Given the one- and two-fermion Green's functions, one can obtain the two-fermion scattering vertex Γ from

$$G_2(p_4, p_3, p_2, p_1) = -\delta_{p_1, p_4} \delta_{p_2, p_3} G(p_1) G(p_2) + \frac{T}{N} \delta_{p_1 + p_2, p_3 + p_4} G(p_4) G(p_3) \Gamma(p_4, p_3, p_2, p_1) G(p_2) G(p_1). \quad (4)$$

With Γ and G one can solve the t -matrix equations represented diagrammatically in Fig. 1 to obtain⁴ the irreducible particle-hole vertex $\bar{\Gamma}_{\text{ph}}(p|p')$ for a center-of-mass momentum $\mathbf{Q}=(\pi, \pi)$ and the irreducible particle-particle vertex $\bar{\Gamma}_{\text{pp}}(p|p')$ for zero center-of-mass momen-

tum. Then, using these, one has the Bethe-Salpeter equations⁵ for the particle-hole channel,

$$-\frac{T}{N} \sum_{p'} \bar{\Gamma}_{\text{ph}}(p|p') G_\uparrow(p'+Q) G_\downarrow(p') \psi_\alpha(p') = \lambda_\alpha \psi_\alpha(p) \quad (5)$$




FIG. 1. The particle-hole and particle-particle t -matrix equations. Here the lines represent dressed single-particle propagators and $\mathbf{Q}=(\pi,\pi)$. Γ is the two-fermion scattering vertex, and $\bar{\Gamma}_{ph}$ and $\bar{\Gamma}_{pp}$ are the irreducible particle-hole and particle-particle vertices, respectively.

and the particle-particle channel,

$$-\frac{T}{N} \sum_{p'} \bar{\Gamma}_{pp}(p|p') G_{\uparrow}(p') G_{\downarrow}(-p') \phi_{\alpha}(p') = \lambda_{\alpha} \phi_{\alpha}(p). \quad (6)$$

Here, as before, the sum on p' sums on both the momentum \mathbf{p}' and the Matsubara frequencies $\omega_{n'}=(2n'+1)\pi T$. The upper limit to the Matsubara frequencies that can be obtained is set by the discrete time interval $\Delta\tau$ used in the Monte Carlo simulation. Here we used $\Delta\tau=0.125$ and a corresponding frequency cutoff of order the bandwidth $8t$.

In Figs. 2(a) and 2(b) we have plotted the leading eigenvalues versus temperature for $U=4t$ and $U=8t$, respectively. These results were obtained for a half-filled 8×8 lattice. If an eigenvalue reaches 1, this implies an instability in the scattering response. At half-filling, the dominant response, the solid circles in Fig. 2, occurs in the particle-hole $\mathbf{Q}=(\pi,\pi)$ channel, reflecting the strong antiferromagnetic fluctuations. The open symbols show the behavior of three of the leading particle-particle eigenvalues of Eq. (6). Comparing the results shown in Figs. 2(a) and 2(b), one can see that increasing U from $4t$ to $8t$ has a dramatic effect on the size of the particle-particle eigenvalues at higher temperatures. For example, at a temperature $T=0.25t$, the $d_{x^2-y^2}$ wave eigenvalue increases by almost a factor of 3 when U is changed from $4t$ to $8t$. In addition, it changes from the second leading eigenvalue to the leading one.

The momentum and frequency dependence of the leading antiferromagnetic eigenvalue, $\psi_s(\mathbf{p}, i\omega_n)$ are shown⁶ in Figs. 3(a) and 3(b) for an 8×8 lattice with $U=8t$ and $T=0.25t$. We see that ψ_s is essentially a constant function of the relative momentum, corresponding to a magnetization operator of the form

$$m_Q^+ = \frac{1}{\sqrt{N}} \sum_{\mathbf{p}} g(\mathbf{p}) c_{\mathbf{p}+\mathbf{Q}\uparrow}^{\dagger} c_{\mathbf{p}\downarrow}, \quad (7)$$

with $g(\mathbf{p}) \simeq 1$. However, it has structure in ω_n implying that retardation plays a role in the antiferromagnetic correlations. Including retardation, the magnetization eigenoperator would have the form





FIG. 2. Eigenvalues of the Bethe-Salpeter equations vs T for (a) $U=4t$ and (b) $U=8t$ for an 8×8 half-filled lattice. The solid points are for the $\mathbf{Q}=(\pi,\pi)$ antiferromagnetic particle-hole eigenvalue of Eq. (3). The open symbols denote the eigenvalues of the $\mathbf{q}=(0,0)$ particle-particle Bethe-Salpeter Eq. (4) with (\circ) corresponding to the singlet $d_{x^2-y^2}$ even-frequency amplitude, (\triangle) to a singlet, odd-frequency p -wave amplitude, and (\square) to a triplet, s -wave odd-frequency amplitude.

$$m_Q^+(\tau) = \int_{-\beta}^{\beta} \frac{d\tau'}{2\beta} \frac{1}{\sqrt{N}} \sum_{\mathbf{p}, i\omega_n} e^{-i\omega_n \tau'} \psi(\mathbf{p}, i\omega_n) \times T c_{\mathbf{p}+\mathbf{Q}\uparrow}^{\dagger}(\tau+\tau') c_{\mathbf{p}\downarrow}(\tau). \quad (8)$$

As seen in Fig. 3(b), $\psi(\mathbf{p}, i\omega_n)$ remains finite over the entire bandwidth implying that the usual magnetization operator $(1/\sqrt{N}) \sum_{\mathbf{p}} c_{\mathbf{p}+\mathbf{Q}\uparrow}^{\dagger} c_{\mathbf{p}\downarrow}$ has a good overlap with the eigenoperator. We also find additional eigenfunctions with smaller eigenvalues. Some of these eigenfunctions are odd in ω_n just as some of the particle-particle eigenfunctions discussed below. However, all of the other eigenvalues are small compared to the dominant antiferromagnetic one.

The momentum and frequency dependence of the leading particle-particle Bethe-Salpeter amplitudes $\phi_d(\mathbf{p}, i\omega_n)$, $\phi_p(\mathbf{p}, i\omega_n)$, and $\phi_s(\mathbf{p}, i\omega_n)$ are shown in Figs. 4(a) and 4(b). At low temperatures, the leading pairing eigenfunction has $d_{x^2-y^2}$ symmetry in momentum space

and is even in frequency. The next leading eigenfunction has p_x (or p_y) symmetry in momentum and is odd in frequency, while the next one has s symmetry in momentum and is also odd in frequency. The p_x (p_y) eigenfunctions correspond to the singlet odd-frequency gap proposed by Balatsky and Abrahams,⁷ while the s -state triplet corresponds to the odd-frequency gap discussed by Brezniskii.⁸ At higher temperatures, the leading eigenvalue in the particle-particle channel corresponds to an odd-frequency s -wave triplet. Comparing Figs. 4(b) and 3(b), one also observes that the pairing eigenvalues decay faster in frequency than the leading magnetic eigenvalue. Thus, as expected, retardation plays a much more important role in the pairing channel. In fact, the odd-frequency amplitudes imply that the equal-time two-fermion part of the corresponding pair-field operator vanishes. Thus retardation is essential for the odd-frequency fluctuations. Note that the effective pair-field eigenoperator has the form of Eq. (8) with $\psi(\mathbf{p}, i\omega_n)$ replaced by $\phi_\alpha(\mathbf{p}, i\omega_n)$ and $c_{\mathbf{p}+\mathbf{Q}\uparrow}^\dagger(\tau+\tau')c_{\mathbf{p}\downarrow}(\tau)$ replaced by $c_{\mathbf{p}\uparrow}^\dagger(\tau+\tau')c_{-\mathbf{p}\downarrow}^\dagger(\tau)$.

We interpret these results as evidence that two holes (or electrons) added to a half-filled 8×8 Hubbard lattice




FIG. 3. Monte Carlo results for the leading magnetic eigenfunction $\psi_s(\mathbf{p}, i\omega_n)$ on an 8×8 lattice with $U=8t$ and $T=0.25t$. (a) Momentum structure of $\psi_s(\mathbf{p}, i\pi T)$. Here \mathbf{p} is taken along the path shown in the inset. (b) Frequency structure of $\psi_s(\mathbf{p}, i\omega_n)$ for $\mathbf{p}=(\pi, 0)$.

lead to a state with strong pairing correlations. The fact that at low temperatures, the pair eigenfunction with the largest eigenvalue has $d_{x^2-y^2}$ symmetry is consistent with previous exact diagonalization⁹ and Monte Carlo studies¹⁰ of 4×4 lattices. In these calculations the ground state of the half-filled 4×4 Hubbard model was found to




FIG. 4. Monte Carlo results for the leading eigenfunctions in the pairing channel, $\phi_d(\mathbf{p}, i\omega_n)$, $\phi_p(\mathbf{p}, i\omega_n)$, and $\phi_s(\mathbf{p}, i\omega_n)$, on an 8×8 lattice with $U=8t$ and $T=0.25t$. (a) Momentum structure of the leading eigenfunction for $\omega_n=\pi T$. Here \mathbf{p} is taken along the path shown in the inset of Fig. 3(a). (b) Frequency structure of the leading eigenfunctions. Here $\phi_d(\mathbf{p}, i\omega_n)$ and $\phi_s(\mathbf{p}, i\omega_n)$ are shown for $\mathbf{p}=(\pi, 0)$, and $\phi_p(\mathbf{p}, i\omega_n)$ is shown for $\mathbf{p}=(\pi/2, \pi/2)$.

have s -wave symmetry, while for $U \gtrsim 3t$, the ground state of the system with two electrons removed (or added) was found to have $d_{x^2-y^2}$ symmetry. Thus the pair-field operator connecting these two states must have $d_{x^2-y^2}$ symmetry. In addition, previous Monte Carlo simulations¹¹ in which pair-field susceptibility with and without the two-particle vertex were calculated, showed that there was an attractive interaction in the $d_{x^2-y^2}$ channel. However, the results reported here differ from previous Monte Carlo calculations in that by studying the Bethe-Salpeter equations, we have allowed the system to select the internal momentum, frequency, and spin structure of the dominant pairing correlations rather than imposing this structure by a particular choice for the pair-field operator. Furthermore, by studying the case where two fermions are added to a half-filled band, we have been

able to run the Monte Carlo simulations at significantly lower temperatures and larger values of U than can be achieved in simulations carried out away from half-filling.

This work was partially supported by the Department of Energy under Grant No. DE-FG03-85ER45197, and by the Electric Power Research Institute under Grant No. RP8009-18. N.B. gratefully acknowledges financial support from IBM. S.R.W. would like to thank the Office of Naval Research for support under Grant No. N00014-91-J-1143. This research was also supported in part by the University of California through an allocation of computer time. The numerical calculations reported in this paper were primarily performed at the San Diego Supercomputer Center.

¹P. W. Anderson and J. R. Schrieffer, Phys. Today 54 (June 1991).

²Starting with a half-filled 8×8 lattice and adding two fermions corresponds to a doping $\delta = 66/64 - 1 \cong 0.03$.

³E. Loh *et al.*, Phys. Rev. B 41, 9301 (1990).

⁴N. Bulut, D. J. Scalapino, and S. R. White, Phys. Rev. B 47, 6157 (1993).

⁵Equivalent forms for these Bethe-Salpeter equations, which directly involve only G and G_2 , can be written as

$$-\sum_p' \frac{G_2(p+Q, p, p', p'+Q)}{G(p+Q)G(p)} \psi_\alpha(p') = \frac{1}{1-\lambda_\alpha} \psi_\alpha(p)$$

and

$$-\sum_p' \frac{G_2(p, -p, -p', p')}{G(p)G(-p)} \phi_\alpha(p') = \frac{1}{1-\lambda_\alpha} \phi_\alpha(p).$$

Note that λ_α , ψ_α , and ϕ_α are the same eigenvalues and eigenfunctions that appear in Eqs. (5) and (6).

⁶The eigenfunctions are normalized such that $(T/N) \sum_p \psi_\alpha^2(p) = 1$.

⁷A. Balatsky and E. Abrahams, Phys. Rev. B 45, 13 125 (1992).

⁸V. L. Berezinskii, Pis'ma Zh. Eksp. Teor. Fiz. 20, 628 (1974) [JETP Lett. 20, 287 (1974)].

⁹A. Parola, S. Sorella, M. Parrinello, and E. Tosatti, Phys. Rev. B 43, 6190 (1991).

¹⁰E. Dagotto, A. Moreo, R. L. Sugar, and D. Toussaint, Phys. Rev. B 41, 811 (1990).

¹¹S. R. White *et al.*, Phys. Rev. B 39, 839 (1989).